

Table 6. Estimated numbers of diagnoses of AIDS in adult or adolescent Hispanics, by diagnosis during 2002, exposure category and place of birth—United States

Exposure category	Place of birth											
	United States		Central/South America		Cuba		Mexico		Puerto Rico		Total ^a	
	No.	%	No.	%	No.	%	No.	%	No.	%	No.	%
Male-to-male sexual contact	1,198	39	376	52	67	50	767	57	292	17	3,202	39
Injection drug use	898	29	82	11	32	24	170	13	788	45	2,235	27
Male-to-male sexual contact and injection drug use	133	4	14	2	7	5	55	4	58	3	299	4
Heterosexual contact	831	27	237	33	28	21	322	24	600	34	2,365	29
Other ^b	43	1	15	2	1	1	30	2	8	0	116	1
Total ^c	3,103	100	724	100	135	100	1,344	100	1,747	100	8,218	100

Note. These numbers do not represent actual cases in persons with a diagnosis of AIDS. Rather, these numbers are point estimates of cases diagnosed that have been adjusted for reporting delays and for redistribution of cases in persons initially reported without an identified risk. The estimates have not been adjusted for incomplete reporting.

^a Includes 189 persons whose place of birth is not among those listed and 976 whose place of birth is unknown.

^b Includes hemophilia, blood transfusion, perinatal, and risk not reported or not identified.

^c Because column totals were calculated independently of the values for the subpopulations, the values in each column may not sum to the column total.